

Received: 2003.03.03
Accepted: 2003.04.04
Published: 2003.05.22

Sound therapy induced relaxation: down regulating stress processes and pathologies

Elliott Salamon¹, Minsun Kim², John Beaulieu³, George B. Stefano¹

¹ Neuroscience Research Institute, State University of New York at Old Westbury, NY 11568, USA

² The Long Island Conservatory, 1125 Willis Avenue, Albertson, NY 11507, USA

³ BioSonic Enterprises, Ltd, P.O. Box 487, High Falls, New York 12440 USA

Summary

The use of music as a means of inducing positive emotions and subsequent relaxation has been studied extensively by researchers. A great deal of this research has centered on the use of music as a means of reducing feelings of anxiety and stress as well as aiding in the relief of numerous pathologies. The precise mechanism responsible for these mediated effects has never been truly determined. In the current report we propose that nitric oxide (NO) is the molecule chiefly responsible for these physiological and psychological relaxing effects. Furthermore this molecules importance extends beyond the mechanistic, and is required for the development of the very process that it mediates. Nitric oxide has been determined to aid in the development of the auditory system and participate in cochlear blood flow. We show that NO is additionally responsible for the induced exhibited physiological effects. We proceed to outline the precise neurochemical pathway leading to these effects. Furthermore we explore the interrelationship between the varying emotion centers within the central nervous system and explain how the introduction of music can mediate its effects via NO coupled to these complex pathways.

key words: music • stress • relaxation • soundtherapy • nitric oxide

Full-text PDF: http://www.MedSciMonit.com/pub/vol_9/no_5/3514.pdf

Word count: 2171

Tables: –

Figures: 1

References: 77

Author's address: George B. Stefano, Neuroscience Research Institute, State University of New York at Old Westbury, P.O.Box 210, Old Westbury, 11568 New York, NY, USA, email: gstefano@sunynri.org

1. DEFINING STRESS

The term 'stress' as defined in the strict biological sense is an event or stimulus that alters the existing homeostasis within a given organism [1]. Some theorists now refer to the 'healthy state' as one of stability in the face of change. Multiple causes of stress add to what is called 'allostatic loading', which can be pathologic if not relieved. The state may be cognitively appraised or non-cognitively perceived. The disturbed organism may either acutely or chronically experience this stimulus. Indeed, the stressor (the stimulus) may even emerge from within the organism itself, such as in interoceptive psychiatric stress. Stress is difficult to define because there are many types of stressors, or stimuli, that can bring on this homeostatic perturbation. Through an extremely complicated homeostatic process, all living organisms maintain their survival in the face of both external and internal 'stressors' [2,3].

Stress when defined as a psychological phenomenon is characterized by feelings of apprehension, nervousness and helplessness, and is commonly present in patients undergoing medical procedures. Past research demonstrates that stress induces numerous types of physiological complications. Stress has been found to cause hypertension, tachycardia and hyperventilation [4], all of which were shown to be linked with ischemia and can cause fluctuations in body temperature, urinary urgency, enlarged pupils, and loss of appetite [5]. Furthermore it has been demonstrated that stress leads to increased cortisol levels, depressing the immune system. Lastly, conditions that arouse stress may actually increase pain [6]. An overwhelming amount of research has been conducted into methods of alleviating the stress response, as well as exploring possible mechanisms by which these methods act.


2. STRESS AND ITS RELATION TO MUSIC

The use of music has consistently been found to reduce stress levels of patients in clinical settings. Mulooly et al. [6] investigated the use of music for postoperative stress and found that patients who underwent an abdominal hysterectomy reported lower stress levels after listening to music when compared to patients who were not exposed to this treatment. Studies [7] have contrasted music to verbal distraction, concluding that although the methods were comparable for the reduction of stress, music was more effective in the reduction of blood pressure. Further studies find [8] that adult patients that listened to music during dialysis were found to have significantly lower blood pressure after their treatment than before. In further studies the effectiveness of music in the reduction of stress has been measured in myocardial infarction patients [8], and in coronary care units [9]. Music has been paired with other therapeutic techniques to reduce stress as well. In a study of pediatric patients, group music therapy sessions, including singing, and instrument playing, were found to decrease observed stress in children before surgery [5]. Guided imagery and music together were found to decrease pain and stress in patients undergoing elective colorectal surgery [4].

3. HOW EMOTIONS CAUSE STRESS AND HOW MUSIC ALLEVIATES IT: CNS PROCESSES

Music and its calming effects have been demonstrated to have a large emotional component. When pleasant music is heard the brains motivation and reward pathways are reinforced with positive emotion mentally linked to the music. This emotionalized memory includes many 'somatic markers', i.e. bodily sensations that accompany emotion and set the feeling tone', feels right' to the person [10]. Clearly, music and the emotion it imparts can be viewed as a process of reinforcing a positive belief so that rational thought can not hinder the strength of the belief (see [11,12]). Indeed, belief in regard to a therapy and/or doctor and/or personal religion, may in fact stimulate physiological processes, enhancing naturally occurring health processes by augmenting their level of performance. Conversely, emotional stresses such as fear and anxiety can induce cardiovascular alterations, such as cardiac arrhythmias [13–15]. These cardiovascular events can be initiated at the level of the cerebral cortex and may involve insular as well as cingulate, amygdalar and hypothalamic processes. Clinically we may see this as elevated cortisol levels and in some instances can induce sudden death in patients with significant coronary artery disease [16]. In addition, heart rate is often altered under stressful conditions. Neurons in the insular cortex, the central nucleus of the amygdala, and the lateral hypothalamus, owing to their role in the integration of emotional and ambient sensory input, may be involved in the emotional link to the cardiovascular phenomenon. These include changes in cardiac autonomic tone with a shift from the cardioprotective effects of parasympathetic predominance to massive cardiac sympathetic activation [13]. This autonomic component, carried out with parasympathetic and sympathetic preganglionic cells via subcortical nuclei from which descending central autonomic pathways arise, may therefore be a major pathway in how belief may affect cardiovascular function. The importance of music and the elicited emotional response (and therefore limbic activation) was further demonstrated in ischemic heart disease when patients with frequent and severe ventricular ectopic rhythms were subjected to psychological stress [13]. The frequency and severity of ventricular ectopic beats increased dramatically during emotional activation of sympathetic mechanisms but not during reflexively-induced increased sympathetic tone.

The hard-wiring of emotion/music and cardiovascular neural systems probably involves many subcortical descending projections from the forebrain and hypothalamus [17–22]. Cardiovascular changes were observed in experiments where the motor cortex surface was stimulated, eliciting tachycardia accompanied by and independent of changes in arterial blood pressure [23]. The 'sigmoid' cortex [23,24], frontal lobe [25–27], especially the medial agranular region [28], subcallosal gyrus [29], septal area [30,31], temporal lobe [32], and cingulate gyrus [32–34] appear to be involved. The insular cortex in cardiac regulation is important because of its high connectivity with the limbic system, suggest-



ing that the insula is involved in cardiac rate and rhythm regulation under emotional stress [35–38].

The amygdala, with respect to autonomic-emotional integration [39,40], is composed of numerous subnuclei, which play a major role in the elaboration of autonomic responses [41]. There are profuse inputs to this region from the insular and orbitofrontal cortices, the parabrachial nucleus, and the nucleus tractus solitarius [42–44]. Amygdalo-tegmental projections are viewed as a critical link in cerebral cortical control of autonomic function [45,46].

The medial hypothalamus is also implicated in cardiac arrhythmogenesis [47]. Beattie and colleagues [47] suggested that hypothalamic projections that descended into the midbrain periaqueductal gray matter, reticular formation, and intermediolateral nucleus of the spinal cord mediate the response. Magoun and colleagues [48] demonstrated that the lateral hypothalamus and wide areas of the lateral tegmentum are also important for autonomic function. The lateral hypothalamus has long been recognized for its role in the regulation of motivation and emotion and the autonomic concomitants of related behaviors [49]. The densest cortical projection to the lateral hypothalamus arises from the infralimbic cortex [50]. Pressor sites within the insular cortex project more heavily to the lateral hypothalamus than do depressor sites and are represented at caudal levels.

Anatomical studies of the lateral hypothalamus demonstrate projections to the periaqueductal gray matter, the parabrachial region, parvocellular formation, dorsal vagal complex, and spinal cord [51,52]. Furthermore, descending projections of the lateral hypothalamus terminate as a capsule around the dorsal motor nucleus of the vagus nerve, which provides secretomotor fibers to the stomach wall, pancreas, and small intestine. These neural patterns might account for the close association of cardiac and gastric responses.

4. NITRIC OXIDE

The very origin of music as a method of stress release has its roots in the early development of the auditory system. In a study by Fessenden and Schacht [53], it was found that the nitric oxide (NO)/cGMP pathway is thoroughly involved in the development and function of the sensory systems, and specifically in the development of the cochlea. Thus NO is involved in the stimulated relaxation from the very development of the organism, to the mechanism by which the relaxation occurs [54]. Cochlear nerve fibers enter the brainstem and are routed through the thalamus to the auditory cortex. It has been demonstrated that it is along this path that the emotion centers within the limbic system are activated (as depicted in Figure 1. the sensation of music enters the diagrammatic neuronal pathway at the limbic system) [55–57]. Furthermore this neuronal pathway from

auditory nerve to cortex was found to be mediated by NO [58].

When we examine NO signaling, we notice two components the constitutive NO synthase (cNOS) endothelial (e) and neuronal (n) isoforms; see [59]. Constitutive NOS (cNOS), as the name implies, is always expressed. When cNOS is stimulated, NO release occurs for a short period of time, but this level of NO can exert profound physiological actions for a longer period of time [59]. NO is not only an immune, vascular and neural signaling molecule, it is also antibacterial [60,61], antiviral [60,61] and it down-regulates endothelial and immunocyte activation and adherence, thus performing vital physiological activities, including vasodilation [59]. Thus NO release subsequent to music listening, has the potential to protect an organism from microbes and physiologic disorders such as hypertension, and also diminishes excessive immune and endothelial activation occurring largely because of vasodilation modulated by NO [59].

The endocannabinoids, anandamide and 2-arachidonoyl glycerol, are naturally occurring cNOS-derived NO-stimulating signaling molecules that are also constitutively expressed [62]. Anandamide, an endogenous endocannabinoid, can also cause NO release from human immune cells, neural tissues and human vascular endothelial cells [63]. Anandamide can also initiate invertebrate immune cell cNOS-derived NO [64]. Estrogen can also stimulate cNOS-derived NO in human immune and vascular cells [63, 65]. We believe that each signaling system performs this common function under different circumstances. Morphine, another naturally occurring animal signal molecule [66], given its long latency before increases in its levels are detected, arises after trauma/inflammation and, through a NO mechanism, down regulates these processes in neural and immune tissues [67]. Anandamide, as part of the ubiquitous arachidonate and eicosanoid signaling cascade, serves to maintain and augment tonal NO in vascular tissues [59]. Estrogen, through NO release, provides an additional pathway by which the system can down-regulate immunocyte and vascular function in women [63]. This may be due to both the immune and vascular trauma associated with cyclic reproductive activities, such as endometrial buildup, when a high degree of vascular and immune activities are occurring. Given the extent of proliferative growth capacity during peak estrogen levels in this cycle, NO may function to enhance down-regulation of the immune system to allow for these changes. Clearly, therefore, enhanced cNOS activity would be a beneficial effect within the concept and time framework of music and the subsequent relaxation it induces. Thus, these signal molecules, especially endocannabinoid and opiate alkaloids [68] have the potential to make you 'feel' good and relax [69], also release NO, which may be a vital part of this complicated process.

5. SIGNALING MOLECULES LEADING TO RELAXATION

As noted above, once individuals undergo a very mild form of work/activity such as music listening, they experience peripheral vasodilation, warming of the skin, a decrease in heart rate and an overwhelming sense of well-being [7,69].

In examining a potential mechanism for the music induced relaxation, besides the over-riding central nervous system output via the autonomic nervous system, peripheral neuro-vascular processes would appear to be important. We surmise NO to be of fundamental importance in this response because of the increase in peripheral temperature, i.e. vasodilation [70]. For a complete review of possible related mechanisms see [59,71-73].

We also surmise, based on current studies, that endothelial derived NO, released through normal pulsations, due to vascular dynamics responding to heart beat [59] as well as ACh stimulated endothelial NO release, may contribute to the effect of NO in inducing smooth muscle relaxation [74]. Furthermore, vascular pulsations may be of sufficient strength to also stimulate nNOS derived NO release, limiting any basal NE actions [74]. Interestingly, nitrosative stress, mediated by involvement of the reactive nitrogen oxide species, N₂O₃ does inhibit dopamine hydroxylase, inhibiting NE synthesis and contributing to the regulation of neurotransmission and vasodilation [75]. This system may provide an autoregulatory mechanism involved in the neuronal control of peripheral vasomotor responses.

6. CONCLUSION

In summary, the music induced relaxation peripherally appears to be mediated by a system of regulation involving NO, as neurotransmitter and as a locally acting hormone. Contingent on the preliminary vasoconstriction and depolarization of the membrane, vasodilation is mediated by NO liberated from vasodilator nerves that activate guanylate cyclase in smooth muscle and produce cGMP. During this stage, NO and NE exist simultaneously. Due to the characteristics of NO, NE no longer mediates vasoconstriction; instead NO activates guanylate cyclase, which produces vasodilation and the relaxation under a depolarized membrane state (see [30,54,76]).

In conclusion, the above findings demonstrate that music has numerous profound effects [4-6,68, see also [77] for social effects] we believe that this occurs via NO, opiate and the above mentioned hormonal system. Furthermore NO has been shown to be a necessary molecule in the development of the auditory system [53], which is required to enable music to act as a relaxant. Taken together we believe that the complex nitric oxide signaling system is the primary and fundamental (from development to mechanism) method by which music acts as a relaxation device.

REFERENCES:

1. McEwen BS: Protective and damaging effects of stress mediation. *New Engl J Med*, 1998; 338: 171-179
2. Chrousos GP, Gold PW: The concepts of stress and stress system disorders: Overview of physical and behavioral homeostasis. *J Am Med Assoc*, 1992; 267
3. Fricchione GL, Stefano GB: The stress response and autoimmunoregulation. *Adv Neuroimmunol*, 1994; 4: 13-28.
4. Tusek DL, Church JM, Strong SA et al: Guided Imagery A Significant Advance in the Care of Patients Undergoing Elective Colorectal Surgery. *Diseases of the Colon and Rectum*, 1996; 40: 172-178
5. Aldridge K: The Use of Music to Relieve Pre-operational Anxiety in Children Attending Day Surgery. *The Australian Journal of Music Therapy*, 1993; 4: 19-35
6. Mullooly VM, Levin RF, Feldman HR: Music for Postoperative Pain and Anxiety. *Journal of New York State Nurses Association*, 1988; 19: 4-7
7. Steelman V: M Intraoperative Music Therapy Effects on Anxiety, Blood Pressure. *AORN Journal*, 1990; 52: 1026-1034
8. Schuster B: L The Effect of Music Listening on Blood Pressure Fluctuations in Adult Hemodialysis Patients. *Journal of Music Therapy*, 3: 146-153
9. Zimmerman LM, Pierson MA, Marker J: 1988. Effects of Music on Patient Anxiety in Coronary Care Units. *Heart and Lung*, 1985; 17: 560-566
10. Damasio AR: *Descartes' Error: Emotion, Reason, and the Human Brain*, Putnam, New York, 1994
11. Stefano GB, Fricchione GL: The biology of deception: Emotion and morphine. *Med Hypotheses* 49, 1995; 51-54
12. Stefano GB, Fricchione GL: The biology of deception: The evolution of cognitive coping as a denial-like process. *Med. Hypotheses*, 1995; 44: 311-314
13. Lown B, DeSilva RA: Roles of psychologic stress and autonomic nervous system changes in provocation of ventricular premature complexes. *Am J Cardiol*, 1978; 41: 979-985
14. Lown B, Verrier RL: Neural activity and ventricular fibrillation. *New Engl J Med*, 1976; 294: 1165-1170
15. Wellens HJ, Vermeulen A, Durrer D: Ventricular fibrillation occurring on arousal from sleep by auditory stimuli. *Circulation*, 1972; 46: 661-665
16. Schiffer F, Hartley LH, Schulman CL, Abelmann WH: Evidence for emotionally-induced coronary arterial spasm in patients with angina pectoris. *Br Heart J*, 1980; 44: 62-66
17. Holstege G: Some anatomical observations on the projections from the hypothalamus to brainstem and spinal cord: an HRP and autoradiographic tracing study in the cat. *J Comp Neurol*, 1987; 260: 98-126
18. Holstege G, Meiners L, Tan K: Projections of the bed nucleus of the stria terminalis to the mesencephalon, pons, and medulla oblongata in the cat. *Exp Brain Res*, 1985; 58: 379-391
19. Hopkins DA: Amygdalotegmental projections in the rat, cat and rhesus monkey. *Neurosci Lett*, 1975; 1: 263-270
20. Hopkins DA, Holstege G: Amygdaloid projections to the mesencephalon, pons and medulla oblongata in the cat. *Exp Brain Res*, 1978; 32: 529-547
21. Kuypers HGJM, Maisky VA: Retrograde axonal transport of horseradish peroxidase from spinal cord to brainstem cell groups in the cat. *Neurosci Lett*, 1975; 1: 9-14
22. Swanson LW, Kuypers HG: The paraventricular nucleus of the hypothalamus: cytoarchitectonic subdivisions and organization of projections to the pituitary, dorsal vagal complex, and spinal cord as demonstrated by retrograde fluorescence double-labeling methods. *J Comp Neurol*, 1980; 194: 555-570
23. Schiff M: Untersuchungen ueber die motorischen Functionen des Grosshirns. *Arch Exp Pathol Pharmacol Naunyn Schmiedeberg*, 1875; 3: 171-179
24. Cerevkov A: Ueber den einfluss der Gehirnhemisphaeren auf das Herz und auf das Gefasssystem, Guseff, Kharkov, Russia, 1892
25. Crouch RL, Thompson JK: Autonomic functions of the cerebral cortex. *J Nerv Ment Dis*, 1939; 89: 328-334
26. Hsu S, Hwang K, Chu H: A study of the cardiovascular changes induced by stimulation of the motor cortex in dogs. *Am J Physiol*, 1942; 137: 468-472
27. Winkler C: Attention and respiration. *Proc Acad Sci Amsterdam*, 1899; 1: 121-138
28. Buchanan SL, Valentine J, Powell DA: Autonomic responses are elicited by electrical stimulation of the medial but not lateral frontal cortex in rabbits. *Behav Brain Res*, 1985; 18: 51-62
29. Hoff EC: The role of the cerebral cortex in the central nervous regulation of cardiovascular function. *Confin Neurol*, 1949; 9: 166-176
30. de la Torre JC, Stefano GB: Evidence that Alzheimer's disease is a microvascular disorder: The role of constitutive nitric oxide. *Brain Res Rev*, 2000; 11: 1581-1585
31. Deutsch DG, Goligorsky MS, Stefano GB et al: Production and physiological actions of anandamide in the vasculature of the rat kidney. *J Clin Invest*, 1997; 100: 1538-1546
32. MacLean PD: Discussion. *Physiol Rev*, 1960; 40: 113-114
33. Smith WK: The functional significance of the rostral cingular cortex as revealed by its responses to electrical excitation. *J Neurophysiol*, 1945; 8: 241-254
34. Ueda H: Arrhythmias produced by cerebral stimulation. *Jpn Circ J*, 1962; 26: 225-230
35. Fimiani C, Liberty T, Acuirre AJ et al: Opiate, cannabinoid, and eicosanoid signaling converges on common intracellular pathways: Nitric oxide coupling. *Prostaglandins*, 1998; 57: 23-34
36. Russchen FT: Amygdalopetal projections in the cat. I. Cortical afferent connections. A study with retrograde and anterograde tracing techniques. *J Comp Neurol*, 1982; 206: 159-179
37. Calaresu FR, Ciriello J: Projections to the hypothalamus from buffer nerves and nucleus tractus solitarius in the cat. *Am J Physiol*, 1980; 239: 130-136
38. Melville KI, Blum G, Shister HE, Silver MD: Cardiac ischemic changes and arrhythmias induced by hypothalamic stimulation. *Am J Cardiol*, 1963; 12: 781-791
39. Davis M: The role of the amygdala in fear and anxiety. *Annu Rev Neurosci*, 1992; 15: 353-375
40. Kapp BS, Frysinger RC, Gallagher M, Haselton JR: Amygdala central nucleus lesions: effect on heart rate conditioning in the rabbit. *Physiol. Behav*, 1979; 23: 1109-1117
41. Mansson E, Bare LA, Yang D: Isolation of a human kappa opioid receptor cDNA from placenta. *Biochem Biophys Res Commun*, 1994; 202: 1431
42. Yasui Y, Itoh K, Kaneko T: Topographical projections from the cerebral cortex to the nucleus of the solitary tract in the cat. *Exp Brain Res*, 1991; 85: 75-84
43. Friedman R, Zuttermeister P, Benson H: Letter to the editor. *New Engl J Med*, 1993; 329: 1201
44. Bonvallet M, Bobo EG: Changes in phrenic activity and heart rate elicited by localized stimulation of amygdala and adjacent structures. *Electroencephalogr. Clin Neurophysiol*, 1972; 32: 1-16
45. Allen GV, Saper CB, Hurley KM, Cechetto DF: Organization of visceral and limbic connections in the insular cortex of the rat. *J Comp Neurol*, 1991; 311: 1-16
46. Kapp BS, Schwaber JS, Driscoll PA: Frontal cortex projections to the amygdaloid central nucleus in the rabbit. *Neuroscience*, 1985; 15: 327-346
47. Beattie J, Brow GR, Long CNH: Physiological and anatomical evidence for the existence of nerve tracts connecting the hypothalamus with spinal sympathetic centres. *Proc R Soc Lond B*, 1930; 106: 253-275
48. Magoun HW, Ranson SW, Heatherington A: Descending connections from the hypothalamus. *Arch Neurol Psychiatry*, 1938; 39: 1127-1149
49. Saper CB, Swanson LW, Cowan WM: An autoradiographic study of the efferent connections of the lateral hypothalamic area in the rat. *J Comp Neurol*, 1979; 183: 689-706
50. Hurley KM, Herbert H, Moga MM, Saper CB: Efferent projections of the infralimbic cortex of the rat. *J Comp Neurol*, 1991; 308: 249-276
51. Hosoya Y, Matsushita M: Brainstem projections from the lateral hypothalamic area in the rat, as studied with autoradiography. *Neurosci Lett*, 1981; 24: 111-116

52. ter Horst GJ, Luiten PG, Kuipers F: Descending pathways from hypothalamus to dorsal motor vagus and ambiguus nuclei in the rat. *J Auton Nerv Syst*, 1984; 11: 59-75
53. Fessenden JD, Schacht J: The nitric oxide/cyclic GMP pathway: a potential major regulator of cochlear physiology. *Hearing Research*, 1998; 118(1-2): 168-176
54. Stefano GB, Goumon Y, Bilfinger TV et al: Basal nitric oxide limits immune, nervous and cardiovascular excitation: Human endothelia express a mu opiate receptor. *Prog Neurobiol*, 2000; 60: 531-544
55. Blood AJ, Zatorre RJ: Intensely pleasurable responses to music correlate with activity in brain regions implicated in reward and emotion. *Proc Natl Acad Sci*, 2001; 25: 98(20): 11818-23
56. Blood AJ, Zatorre RJ, Bermudez P, Evans AC: Emotional responses to pleasant and unpleasant music correlate with activity in paralimbic brain regions. *Nat Neurosci*, 1999; 2(4): 382-7
57. Zatorre RJ, Evans AC, Meyer E: Neural mechanisms underlying melodic perception and memory for pitch. *J Neurosci*, 1994; 14(4): 1908-19
58. Michel O, Hess A, Bloch W et al: Localization of the NO/cGMP-pathway in the cochlea of guinea pigs. *Hear Res*, 1999; 133(1-2): 1-9
59. Chung JW, Schacht J: ATP and nitric oxide modulate intracellular calcium in isolated pillar cells of the guinea pig cochlea. *J Assoc Res Otolaryngol*, 2001; 2(4): 399-407
60. Benz D, Cadet P, Mantione K et al: (2002) Tonal nitric oxide and health: A free radical and a scavenger of free radicals. *Med Sci Monitor*, 2002; 8: 1-4
61. Benz D, Cadet P, Mantione K et al: Tonal nitric oxide and health: Anti-Bacterial and -Viral Actions and Implications for HIV. *Med Sci Monitor*, 2002; 8(1): 27-31
62. Stefano GB, Salzet M, Magazine HI, Bilfinger: Antagonism of LPS and IFN- γ ! Nieznany argument przełącznika. induction of iNOS in human saphenous vein endothelium by morphine and anandamide by nitric oxide inhibition of adenylate cyclase. *J Cardiovasc Pharmacol*, 1998; 31: 813-820
63. Stefano GB, Salzet M, Bilfinger: Long-term exposure of human blood vessels to HIV gp120, morphine and anandamide increases endothelial adhesion of monocytes: Uncoupling of nitric oxide. *J Cardiovasc Pharmacol*, 1998; 31: 862-868
64. Stefano GB, Salzet B, Salzet M: Identification and characterization of the leech CNS cannabinoid receptor: Coupling to nitric oxide release. *Brain Res*, 1997; 753: 219-224
65. Stefano GB, Cadet P, Breton C et al: Estradiol-stimulated nitric oxide release in human granulocytes is dependent on intracellular calcium transients: Evidence for a cell surface estrogen receptor. *Blood*, 2000; 95: 3951-3958
66. Stefano GB, Goumon Y, Casares F et al: Endogenous morphine. *Trends Neurosci*, 2000; 9: 436-442
67. Lembo G, Vecchione C, Izzo R et al: Noradrenergic vascular hyper-responsiveness in human hypertension is dependent on oxygen free radical impairment of nitric oxide activity. *Circulation*, 2000; 102: 552-557
68. Tonnesen E, Brix-Christensen V, Bilfinger TV et al: Endogenous morphine levels increase following cardiac surgery: Decreasing proinflammatory cytokine levels and immunocyte activity. *Int J Cardiol*, 1998; 62: 191-197
69. Snyder M, Chlan L: Music Therapy. *Annu Rev Nurs Res*, 1999; 17: 3-25
70. Krumhansl CL: An exploratory study of musical emotions and psychophysiology. *Can J Exp Psychol*, 1997; 51(4): 336-53
71. Stuart EM, Caudill J, Leserman C et al: Nonpharmacologic treatment of hypertension: A multiple-risk-factor approach. *J Cardiovasc Nurs*, 1987; 1: 1-14
72. Okamura T, Ayajiki K, Uchiyama M et al: Neurogenic vasodilation of canine isolated small labial arteries. *J Pharmacol Exp Ther*, 1999; 288: 1031-1036
73. Toda N: Mediation by nitric oxide of neurally-induced human cerebral artery relaxation. *Experientia*, 1993; 49: 51-53